Geranium Plant Named 'Fisrohot'

Genus and species of the plant claimed:

Hybrid of Pelargonium zonale L'Héritier

Variety denomination:

'Fisrohot'

5

10

15

Background of the Invention

The present invention comprises a new and distinct cultivar of geranium, botanically known as *Pelargonium zonale*, and hereinafter referred to by the cultivar name 'Fisrohot'.

'Fisrohot' is a product of a planned breeding program which had the objective of creating new zonal geranium cultivars with semi-double flowers, relatively vigorous, but well-branched growth habit, good outdoor performance, and in various flower colors.

'Fisrohot' originated from a hybridization made by the inventor, Angelika Utecht, in a controlled breeding program in Hillscheid, Germany, in 1998. The female parent was an unpatented hybrid seedling, no. 97-2191-1, with large, violet, semi-double flowers, dark green foliage with only slight zonation, and medium to vigorous growth habit. The male parent of 'Fisrohot' was the unpatented hybrid seedling no. 97-2001-3, which was derived from self-pollinating variety 'Fislypso' (U.S. Plant Patent no. 11,162), and characterized by deep violet, semi-double flowers, medium to light green foliage without zonation, and low plant habit.

'Fisrohot' was selected as one flowering plant within the progeny of the stated cross by Angelika Utecht in 1999 in a controlled environment in Moncarapacho,

Portugal.

25

20

The first act of asexual reproduction of 'Fisrohot' was accomplished when vegetative cuttings were taken from the initial selection in the fall of 1999, in a controlled environment in Moncarapacho, Portugal, by, or under the supervision of, Angelika Utecht.

Horticultural examination of plants grown from cuttings of the plant initiated in May 2000, in Hillscheid, Federal Republic of Germany, and continuing thereafter, has demonstrated that the combination of characteristics as herein disclosed for 'Fisrohot' are firmly fixed and are retained through successive generations of asexual reproduction.

5

10

15

20

'Fisrohot' has not been observed under all possible environmental conditions.

The phenotype may vary significantly with variations in environment such as temperature, light intensity and day length.

Brief Summary of the Invention

The following observations, measurements, and comparisons describe plants grown in Hillscheid, Germany, under greenhouse conditions which approximate those generally used in commercial practice. The following traits have been repeatedly observed and are determined to be basic characteristics of 'Fisrohot' in combination distinguish this geranium as a new and distinct cultivar:

- 1. Brilliant purple-violet flowers with small, white eyes;
- 2. Large, round inflorescences, long strong peduncles;
- 3. Large, medium-green leaves, with weak zonation;
- 4. Vigorous growth, tall and well-branched plant habit; and
- 5. Medium to late spring flowering response.

Inventor: Anglika Utecht

Of the many commercial cultivars known to the present inventor, the most

similar in comparison to 'Fisrohot' are the varieties 'Fip 750' (U.S. Plant Patent no.

14,011), and 'Fisrovio' (U.S. Plant Patent no. 13,250).

In comparison to 'Fip 750', 'Fisrohot' has a similar main flower color, but

medium-green, instead of dark-green, foliage, a much taller plant habit, and later

begin of flowering.

In comparison to 'Fisrovio', flowers of 'Fisrohot' have a somewhat more

bluish hue, they lack the orange eyes on petals, but show white bases instead, the

zonation on leaves is somewhat more distinct, and internodes of branches are longer.

10

5

Brief Description of the Drawing

The accompanying photographic drawing shows typical flower and foliage

characteristics of 'Fisrohot' with colors being as true as possible with an illustration

of this type. The photographic drawing shows a flowering potted 'Fisrohot' plant.

15

20

Detailed Botanical Desription

The measurements were taken in Hillscheid, Germany, in mid May 2003, 11

weeks after planting of rooted cuttings. The plants were growing in 14 cm pots, they

had not been pinched. In the following description color references are made to the

Royal Horticultural Society Color Chart. The color values were determined indoors

from plants growing in a green-house in May 2003 in Hillscheid, Germany.

INFLORESCENCE

Umbel:

Shape:

Semi-spherical to nearly spherical

25 Average diameter:

117 mm

3

Average depth:

65-70 mm

Peduncle:

233 mm in length, 4 - 4.5 mm in diameter

Peduncle color:

Light green, RHS 143 C, one side may show a tinge of

brown RHS 173 A

5

Pedicel:

31 mm in length

Pedicel color:

Light green, RHS 143 B, partly brownish infused, RHS

179 A

Number of flowers per umbel:

About 45-65

Corolla:

10 Average diameter:

53 mm

Form:

Semi-double type

Shape:

Round outline, flat cup-shape, with the upper petals

about the same size as the lower petals

Number of petals:

Approximately 7-9

15

Number of petaloids:

None

Shape of petals:

Obovate, base acute, upper end is truncate or rounded,

margin is entire or very weakly crenated, and often

wavy

Size of petals:

Upper petals: 26-27 mm long, 21-23 mm wide;

20

lower petals: 25-26 mm long, 22-23 mm wide

Color (general tonality from a distance of three meters):

Bright violet purple

Color of upper petals:

Main part between RHS N74 A and 74 B

Markings of upper petals:

White bases more then a third of the petal, with

fine pink veins RHS 68 B

25 Color of lower petals:

RHS 74 B, uniform

Markings of lower petals: None

Color of lower surface of petals: RHS 67 B

Color of sepals: Outer surface: light green, RHS 144 A; inner surface:

light green, RHS 144 B

5 Number of sepals: 5

Shape of sepals: linear to lanceolate, acute tip, truncate base, surface with

moderate pubescence, margin entire

Size of sepals: 12-14 mm long, 4 mm wide for the largest upper sepal,

2-3 mm in width for the other sepals

10 Bud: (just prior to petals unfolding)

Shape: Elliptical

Color of sepals: Light green, RHS 143 C

Color of petals: RHS 68 B

Length: 20-21 mm

15 Width: 12-13 mm

REPRODUCTIVE ORGANS:

Androecium: 5-7 fertile anthers, plenty pollen, yellow-orange, RHS 26 A,

filaments white, RHS 155 D, to light-pink, RHS 52 D

20 Gynoecium: One pistil, whitish style, RHS 155 D to 65 D, stigma 5- 6-lobed

stigma, dark-red, RHS 53 A

Fertility/seed seed: No seed set observed

Spring flowering response period: In Hillscheid, Germany, in 2001 plants had on

average .3 flowers opened 8 weeks after planting

of rooted cuttings

Outdoor flower production: Continuously and rich flowering, the flower

count in 2003, in Hillscheid, Germany, indicated

about 1-2 inflorescence per plant in mid May

Durability: Good stability of flower color, good rain resistance

Lastingness of the individual flower: About 8 days at 18°C, about 16 days for

the umbel

10 Fragrance: None

PLANT

5

Foliage:

Shape:

Kidney-shaped, with cordate base, with the gap between

the lowest lobes open, apex rounded with weak lobes

Margin: Bi-crenated

Texture: Upper surface smooth, dull

Size of leaf: 115 mm wide, 75 mm long

Color of upper surface: Medium green, approximately RHS 137 C

20 Color of zonation: Brown, about RHS 166 A, distinctness weak to

medium

Color of lower surface: RHS 143 B

Petioles: 90 mm long, 2-2.5 mm diameter, green in color, approximately

RHS 137 D

15

General appearance and form:

Stem color:

Light green, RHS 143 C

Internode length:

30-40 mm

Branching pattern:

4-6 branches

5 Size of plants:

20.3 cm tall, 37.1 cm wide, (11 -week-old plants, as

described, measured from the top of the soil (base of the

main stem) to the surface of the foliage canopy, without

inflorescences